

## Process for preparing optically pure active compounds

### Subject-matter of the invention

The present invention relates to a novel process for preparing optically pure active compounds which can be used for preparing medicaments in the pharmaceutical industry.

### Technical background

Pyridin-2-ylmethylsulphonyl-1H-benzimidazoles and compounds of a closely related structure, as known, for example, from EP-A-0005129, EP-A-0166287, EP-A-0174726 and EP-A-0268956, are, owing to their  $H^+/K^+$ -ATPase-inhibitory action, of considerable importance in the therapy of diseases associated with an increased secretion of gastric acid.

Examples of active compounds from this class of compounds which are commercially available or in clinical development are 5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulphonyl]-1H-benzimidazole (INN: omeprazole), (S)-5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulphonyl]-1H-benzimidazole (INN: esomeprazole), 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole (INN: pantoprazole), 2-[3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridinyl)methylsulphonyl]-1H-benzimidazole (INN: lansoprazole), 2-[[4-(3-methoxypropoxy)-3-methylpyridin-2-yl)methylsulphonyl]-1H-benzimidazole (INN: rabeprazole) and 5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridylmethyl)sulphonyl]-1H-imidazo(4,5-b)pyridine (INN: tenatoprazole).

The abovementioned sulphonyl derivatives which, owing to their mechanism of action, are also referred to as proton pump inhibitors or abbreviated PPI are chiral compounds. The process usually used for preparing the PPI is the oxidation of the corresponding sulphides. This oxidation gives – unless particular measures are taken – a racemic mixture comprising about the same proportions of the two enantiomers (stereoisomers), i.e. the (+)- and (-)-form or the (R)- and (S)-form of the PPI.

Since enantiomers are thermally relatively stable, i.e. they do not racemize on storage – in particular in solid form – there has in the past been no lack of efforts to separate PPI enantiomer mixtures or to prepare the PPI enantiomers in more or less pure form.

### Prior art

The international patent application WO91/12221 describes a process for separating enantiomers using a cellulase enzyme. One of the active compounds mentioned as being separable into the enantiomers with the aid of this process is omeprazole.

The international patent application WO92/08716 describes, for the first time, a chemical process which allows the separation of pyridin-2-ylmethylsulphonyl-1H-benzimidazoles into their optical isomers.

10/536766  
536,766

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
24 June 2004 (24.06.2004)

PCT

(10) International Publication Number  
**WO 2004/052882 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 401/12**,  
471/04

(81) Designated States (*national*): AE, AL, AU, BA, BR, CA, CN, CO, DZ, EC, EG, GE, HR, ID, IL, IN, IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, SG, TN, UA, US, VN, YU, ZA, ZW.

(21) International Application Number:  
PCT/EP2003/013605

(22) International Filing Date: 3 December 2003 (03.12.2003)

(84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
02027273.8 6 December 2002 (06.12.2002) EP  
103 40 255.1 29 August 2003 (29.08.2003) DE

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AL, AU, BA, BR, CA, CN, CO, DZ, EC, EG, GE, HR, ID, IL, IN, IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, SG, TN, UA, VN, YU, ZA, ZW, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR)
- of inventorship (Rule 4.17(iv)) for US only

(71) Applicant (*for all designated States except US*): **ALTANA PHARMA AG** [DE/DE]; Byk-Gulden-Str. 2, 78467 Konstanz (DE).

**Published:**

- with international search-report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **KOHL, Bernhard** [DE/DE]; Zum Brühl 9, 78465 Konstanz (DE). **MÜLLER, Bernd** [DE/DE]; Bücklestr. 84a, 78467 Konstanz (DE). **WEINGART, Ralf Steffen** [DE/DE]; Thingoltstr. 34, 78465 Konstanz (DE).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(74) Agent: **WOLF, Ulrich**; Altana Pharma AG, Byk-Gulden-Str.2, 78467 Konstanz (DE).

(54) Title: **PROCESS FOR PREPARING OPTICALLY PURE ACTIVE COMPOUNDS**

(57) Abstract: The invention relates to a novel process for preparing optically pure PPI having a sulphinyl structure using a chiral zirconium complex or a chiral hafnium complex.

WO 2004/052882 A1

Compounds mentioned as having been prepared in an exemplary manner are, inter alia, the compounds (+)- and (-)-5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole [= (+)- and (-)-pantoprazole]. The international patent application WO92/08716 refers to the fact that the optical isomers of the pyridin-2-yl-methylsulphonyl-1H-benzimidazoles, i.e. the (+)- and (-)-enantiomers or the (R)- and (S)-enantiomers, are used as active compounds in medicaments for the treatment of gastrointestinal disorders. With respect to the mode of application and the dosage of the active compounds, reference is made inter alia to the European patent 166 287.

The international patent application WO94/27988 describes the separation of racemic omeprazole into the enantiomers, using chiral auxiliaries.

The international patent application WO96/02535 (= USP 5,948,789) describes a process for the enantioselective synthesis of PPI using chiral titanium complexes. What is described is, inter alia, the synthesis of (+)- and (-)- [or, expressed in a different way, (R)- and (S)]-pantoprazole, the chiral auxiliary used for the synthesis of (+)-pantoprazole being diethyl (+)-tartrate and the chiral auxiliary used for the preparation of (-)-pantoprazole being diethyl (-)-tartrate.

The international patent applications WO96/17076 and WO96/17077 describe the enantioselective biooxidation or bioreduction with the use of certain microorganisms for the preparation of enantiomerically pure or enantiomerically enriched PPI.

The international patent application WO97/02261 describes the enrichment of PPI enantiomers by selective precipitation.

The international patent applications WO94/24867 and WO94/25028 claim the use of the compounds (-)- and (+)-pantoprazole for treating stomach disorders in humans. Each of the stereoisomers is said to have medical advantages compared to the respective other stereoisomers.

The enantioselective sulphoxidation for preparing esomeprazole ((S)-omeprazole) on a large scale using a chiral titanium complex is described in *Tetrahedron, Asymmetry*, (2000), 11, 3819-3825.

The enantioselective sulphoxidation of aryl alkyl sulphides and dialkyl sulphides in the presence of a zirconium catalyst having a polydentate ligand is described in *J. Org. Chem.*, (1999), 64(4), 1327.

### **Description of the invention**

The invention provides a process for preparing optically pure PPI having a sulphonyl structure. The process is characterized in that the oxidation of the corresponding sulphide is carried out in the presence of a chiral zirconium complex or a chiral hafnium complex, the chiral auxiliary used being an optically pure tartaric acid derivative.

The oxidation is advantageously carried out in an organic solvent, such as, for example, ethyl acetate, toluene, dichloromethane, dioxane or, preferably, methyl isobutyl ketone, where it is not necessary for the solvents mentioned to be completely anhydrous or where anhydrous solvents are in each case optionally admixed with a defined proportion of water, for example up to a maximum of 0.5 equivalent. For reactions with less than 0.5 equivalent of zirconium or hafnium complex, it is preferred to use an anhydrous solvent. The solvents employed may be used in the commercially available quality.

A solvent essentially comprises a specific solvent if it contains at least 50%, preferably at least 90%, in particular at least 95%, of said specific solvent. An anhydrous solvent is essentially free of water, having a water content of less than 5%, preferably less than 1%, in particular less than 0.3%.

Suitable oxidizing agents are all anhydrous oxidizing agents customarily used for the synthesis of PPI, where particular mention may be made of hydroperoxides, such as, for example, tert-butyl hydroperoxide or, in particular, cumene hydroperoxide. In general, 0.90 to 1.3 oxidation equivalents, preferably 0.95-1.05 equivalents, of the oxidizing agent are used.

Suitable zirconium complexes are, for example, zirconium(IV) acetylacetonate, zirconium(IV) butoxide, zirconium(IV) tert-butoxide, zirconium(IV) ethoxide and, in particular, zirconium(IV) n-propoxide (preferably as a solution in n-propanol) or zirconium(IV) isopropoxide (preferably in the form of the zirconium(IV) isopropoxide/isopropanol complex). Suitable hafnium complexes are, for example, hafnium(IV) acetylacetonate, hafnium(IV) butoxide, hafnium(IV) n-propoxide, hafnium(IV) isopropoxide (preferably in the form of the hafnium(IV) isopropoxide/isopropanol complex), hafnium(IV) ethoxide and in particular hafnium(IV) tert-butoxide. Preference is given to using a zirconium complex.

In general, 0.01-2 equivalents, preferably 0.05-0.9 equivalent, of the zirconium complex or of the hafnium complex are used.

Suitable optically pure tartaric acid derivatives are, for example (+)-L-tartaric acid amides, such as (+)-L-tartaric acid bis-(N,N-diallylamide), (+)-L-tartaric acid bis-(N,N-dibenzylamide), (+)-L-tartaric acid bis-(N,N-diisopropylamide), (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-piperidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), (+)-L-tartaric acid bis-(N-cycloheptylamide) or (+)-L-tartaric acid bis-(N-4-methyl-N-piperazinamide), or dialkyl (+)-L-tartrates, such as dibutyl (+)-L-tartrate, di-tert-butyl (+)-L-tartrate, diisopropyl (+)-L-tartrate, dimethyl (+)-L-tartrate and diethyl (+)-L-tartrate, or (-)-D-tartaric acid amides, such as (-)-D-tartaric acid bis-(N,N-diallylamide), (-)-D-tartaric acid bis-(N,N-dibenzylamide), (-)-D-tartaric acid bis-(N,N-diisopropylamide), (-)-D-tartaric acid bis-(N,N-dimethylamide), (-)-D-tartaric acid bis-(N-pyrrolidinamide), (-)-D-tartaric acid bis-(N-piperidinamide), (-)-D-tartaric acid bis-(N-morpholinamide), (-)-D-tartaric acid bis-(N-cycloheptylamide) or (-)-D-tartaric acid bis-(N-4-methyl-N-piperazinamide), or

dialkyl (-)-D-tartrates, such as dibutyl (-)-D-tartrate, di-tert-butyl (-)-D-tartrate, diisopropyl (-)-D-tartrate, dimethyl (-)-D-tartrate and diethyl (-)-D-tartrate. In general, 0.02-4 equivalents, preferably 0.1-2 equivalents, of the optically pure tartaric acid derivative are employed.

Particularly preferred (+)-L-tartaric acid derivatives are (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), and particularly preferred (-)-D-tartaric acid derivatives are (-)-D-tartaric acid bis-(N,N-dimethylamide), (-)-D-tartaric acid bis-(N-pyrrolidinamide), (-)-D-tartaric acid bis-(N-morpholinamide).

Particularly suitable for the preparation of (-)-pantoprazole are (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide) and (+)-L-tartaric acid bis-(N-morpholinamide), and particularly suitable for the preparation of (+)-pantoprazole are (-)-D-tartaric acid bis-(N,N-dimethylamide), (-)-D-tartaric acid bis-(N-pyrrolidinamide) and (-)-D-tartaric acid bis-(N-morpholinamide).

The oxidation is preferably carried out at temperatures between -20 and 50°C, in particular at room temperature and optionally in the presence of a base, suitable bases being, in particular, organic bases, preferably a tertiary amine, such as triethylamine or N-ethyldiisopropylamine.

If the process is carried out in a suitable manner, the optically pure PPI having sulphonyl structure is obtained in an optical purity of >95%. By further steps, such as, for example, pH-controlled reprecipitation and/or recrystallization in a suitable solvent, such as, for example, isopropanol, it is possible to further increase the optical purity considerably. Reprecipitation is carried out via intermediate preparation of suitable salts, such as, for example, via the sodium salt (for other possible salts, see, for example, EP-A-166287).

The invention is illustrated in more detail by the examples below, but not limited in any way. The abbreviation h stands for hour(s).

### Examples

**1. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole  
[= (-)-pantoprazole or (S)-pantoprazole] with diethyl (+)-L-tartrate and zirconium(IV)  
isopropoxide/isopropanol**

A) At room temperature, 20.2 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole together with 17.9 g of diethyl (+)-tartrate, 13.4 g of zirconium(IV) isopropoxide/isopropanol and 0.1 ml of water are suspended in 100 ml of methyl isobutyl ketone. The mixture is heated at 40°C for one hour, resulting in the formation of an almost clear solution. After cooling to room temperature, 4.1 ml of N-ethyldiisopropylamine are added. 11 ml of cumene hydroperoxide are then slowly metered in. Stirring at room temperature is continued until the oxidation process has ended (monitored by TLC). The clear solution is quenched with 0.9 g of sodium thiosulphate in 54 ml of water and 30.3 g of 40% (w/w) of NaOH and stirred for another 14 h. After addition of 25 g of sodium chloride, the phases are separated. The aqueous phase is extracted with 50 ml of methyl isobutyl ketone. The combined organic phases are washed together using 25 ml of saturated sodium chloride solution. 150 ml of water are added to the methyl isobutyl ketone solution, and the pH is adjusted to 13 using 10% (w/w) NaOH. The phases are separated and the methyl isobutyl ketone phase is extracted once more with 50 ml of water at pH 13. The aqueous phases are combined and, at 40°C and under reduced pressure, subjected to incipient distillation. At 40-50°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH 9. Under pH control, stirring is continued for another 12 h. The beige crystals are filtered off and washed with 50 ml of water. This gives the title compound in an optical purity of >90%.

To increase the purity, (-)-pantoprazole is dissolved in water/NaOH and again precipitated by addition of acetic acid to pH 9. Drying gives a beige powder of melting point 145°C (decomposition) and an optical purity of >95%. If this powder is recrystallized from 2-PrOH, a clear crystal of melting point 147-149°C (decomposition) with an optical rotation of  $\alpha_D^{20} = -140$  (c=0.5, MeOH) is obtained.

B) Alternatively, the reaction described in Example 1A can be carried out in 100 ml of toluene instead of methyl isobutyl ketone. If the reaction is carried out in toluene, the zirconium salts have to be filtered off after quenching and the reaction product ((S)-pantoprazole as sodium salt) is directly extracted into the aqueous phase. From the aqueous phase, it can then be precipitated under controlled pH as (S)-pantoprazole. This gives beige crystals of an optical purity of > 95%.

2. **(-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole**  
**[ = (-)-pantoprazole or (S)-pantoprazole] with (+)-L-tartaric acid bis-(N,N-dimethylamide)**  
**and zirconium(IV) isopropoxide/isopropanol**

At room temperature, 20.2 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole are suspended in 100 ml of methyl isobutyl ketone together with 18.0 g of (+)-L-tartaric acid bis-(N,N-dimethylamide) and 13.4 g of zirconium(IV) isopropoxide-isopropanol. The mixture is heated at 40°C for one hour, resulting in the formation of a solution which is almost clear. After cooling to room temperature, 4.1 ml of N-ethyl-diisopropylamine are added. 11 ml of cumene hydroperoxide are then slowly metered in. Stirring is continued at room temperature until the oxidation has ended (5-10 hours, monitored by TLC). The clear solution is diluted with 100 ml of methyl isobutyl ketone and quenched with 1.8 g of sodium thiosulphate in 140 ml of water and stirred for a further 14 hours. After phase separation, 55 ml of saturated sodium bicarbonate solution and 55 ml of methyl isobutyl ketone are added to the aqueous phase, and the phases are separated. Another 55 ml of saturated sodium bicarbonate solution and 55 ml of methyl isobutyl ketone are added to the aqueous phase, and the phases are separated. The combined methyl isobutyl ketone phases are then washed twice with 55 ml of saturated sodium bicarbonate solution. 150 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to pH = 13 using a 40% by weight strength aqueous solution of sodium hydroxide. After phase separation, the methyl isobutyl ketone phase is extracted with another 50 ml of water at pH = 13. The aqueous phases are combined and, at 40°C, subjected to incipient distillation under reduced pressure. At 40-45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH = 9.0. Stirring is continued for another 12 hours during which the pH is monitored. The beige crystals are filtered off and washed with 50 ml of water. The title compound is obtained in a yield of about 15 g (73% of theory) and an optical purity of >95%.

To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and re-precipitated with acetic acid (10%) at pH = 9.0.

3. **(-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole**  
**[ = (-)-pantoprazole or (S)-pantoprazole] with (+)-L-tartaric acid bis-(N,N-pyrrolidinamide)**  
**and zirconium(IV) isopropoxide/isopropanol**

At room temperature, 20.2 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole are suspended in 100 ml of methyl isobutyl ketone together with 22.6 g of (2R,3R)-(+)-L-tartaric acid bis-(N-pyrrolidinamide) and 13.4 g of zirconium(IV) isopropoxide-isopropanol. The mixture is heated at 40°C for one hour, resulting in the formation of a solution which is almost clear. After cooling to room temperature, 4.1 ml of N-ethyl-diisopropylamine are added. 11 ml of cumene hydroperoxide are then slowly metered in. Stirring is continued at room temperature until the oxidation has ended (5-10 hours, monitored by TLC). The clear solution is diluted with 100 ml of methyl isobutyl ketone and quenched with 1.8 g of sodium thiosulphate in 140 ml of saturated sodium bicarbonate solution and stirred for a further 14 hours. After phase separation, the mixture is washed twice with

55 ml of saturated sodium bicarbonate solution. 150 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to pH = 13 using a 40% by weight strength aqueous solution of sodium hydroxide. After phase separation, the methyl isobutyl ketone phase is extracted with another 50 ml of water at pH = 13. The aqueous phases are combined and, at 40°C, subjected to incipient distillation under reduced pressure. At 40-45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH = 9.0. Stirring is continued for another 12 hours during which the pH is monitored. The beige crystals are filtered off and washed with 50 ml of water. The title compound is obtained in a yield of about 17 g (80% of theory) and an optical purity of >98%.

To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and re-precipitated with acetic acid (10%) at pH = 9.0.

**4. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with (+)-L-tartaric acid bis-(N,N-pyrrolidinamide) and zirconium(IV) n-propoxide**

At room temperature, 20.2 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole are suspended in 100 ml of methyl isobutyl ketone together with 22.6 g of (+)-L-tartaric acid bis-(N-pyrrolidinamide) and 16.5 g of zirconium(IV) n-propoxide (70% in propanol). The mixture is heated at 40°C for one hour, resulting in the formation of a solution which is almost clear. After cooling to room temperature, 4.1 ml of N-ethyl-diisopropylamine are added. 10 ml of cumene hydroperoxide are then slowly metered in. Stirring is continued at room temperature until the oxidation has ended (5-24 hours, monitored by TLC). The clear solution is diluted with 100 ml of methyl isobutyl ketone and quenched with 1.8 g of sodium thiosulphate in 140 ml of saturated sodium bicarbonate solution and stirred for a further 14 hours. After phase separation, the mixture is washed twice with 55 ml of saturated sodium bicarbonate solution. 150 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to pH = 13 using a 40% by weight strength aqueous solution of sodium hydroxide. After phase separation, the methyl isobutyl ketone phase is extracted with another 50 ml of water at pH = 13. The aqueous phases are combined and, at 40°C, subjected to incipient distillation under reduced pressure. At 40-45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH = 9.0. Stirring is continued for another 12 hours during which the pH is monitored. The beige crystals are filtered off and washed with 50 ml of water. The title compound is obtained in a yield of about 16 g (75% of theory) and an optical purity of >98%.

To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and re-precipitated with acetic acid (10%) at pH = 9.0.



5. **(-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole**  
**[ = (-)-pantoprazole or (S)-pantoprazole] with (+)-L-tartaric acid bis-(N,N-pyrrolidinamide)**  
**and zirconium(IV) n-propoxide**

Analogously to Example 4, reaction of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole under otherwise identical conditions, but without addition of N-ethyl-diisopropylamine, gives the title compound in a yield of 65% of theory and an optical purity of >98%.

6. **(-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole**  
**[ = (-)-pantoprazole or (S)-pantoprazole] with catalytic amounts of (+)-L-tartaric acid bis-**  
**(N-pyrrolidinamide) and zirconium(IV) n-propoxide**

Analogously to Example 4, reaction of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole under otherwise identical conditions, but with 0.1 equivalent of zirconium n-propoxide, 0.25 equivalent of (+)-L-tartaric acid bis-(N-pyrrolidinamide) and 0.07 equivalents of Hünig base gives, after an oxidation time of 48-72 h, the title compound in a yield of 80% of theory and an optical purity of >98%.

7. **(-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole**  
**[ = (-)-pantoprazole or (S)-pantoprazole] with catalytic amounts of (+)-L-tartaric acid bis-**  
**(N-pyrrolidinamide) and zirconium(IV) n-propoxide**

At room temperature, 50.0 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole and 5.2 g of (+)-L-tartaric acid bis-(N-pyrrolidinamide) (0.15 eq.) are suspended in 360 ml of methyl isobutyl ketone (MIBK). The suspension is heated at 40-45°C and 60 ml of MIBK are distilled off for azeotropic removal of water present in the mixture. At this temperature, 3.2 g of zirconium(IV) n-propoxide (70% in propanol, 0.05 eq.) are added, and stirring is continued for 1 hour. After cooling to 30°C, 0.9 ml of N-ethyl-diisopropylamine are added. 27.1 g of cumene hydroperoxide (80% in cumene) are then slowly metered in. Stirring is continued at 30°C until the exothermic oxidation process has ended (20 hours, monitored by TLC or HPLC). The suspension is diluted with 60 ml of 2-propanol and quenched with 1.69 g of sodium thiosulphate in 100 ml of saturated sodium bicarbonate solution and stirred for at least 2 hours. After phase separation, the mixture is washed twice with 50 ml of saturated sodium bicarbonate solution. 150 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to 12.5-13 using 10 ml of aqueous sodium hydroxide solution (40% (w/w)). After phase separation, the methyl isobutyl ketone phase is extracted 2 more times with 100 ml of water and 2 ml of aqueous sodium hydroxide solution (40% (w/w)) at pH = 12.5-13. The combined aqueous phases are reextracted twice with 50 ml of methyl isobutyl ketone and subjected to incipient distillation at 40°C under reduced pressure. At 40-45°C, (-)-pantoprazole is

precipitated by addition of 10% strength acetic acid to pH = 9.0. Under pH control, stirring is continued for another 12 hours. The beige crystals are filtered off and washed twice with 50 ml of water. This gives the title compound in a yield of 82% of theory in a chemical purity of 95% and an optical purity of > 95%.

To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and again precipitated at pH = 9.0 using acetic acid (10%). This gives the title compound in a yield of 75% of theory in a chemical purity of > 97% and an optical purity of > 98%.

**8. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole  
[= (-)-pantoprazole or (S)-pantoprazole] with catalytic amounts of (+)-L-tartaric acid bis-  
(N-pyrrolidinamide) and zirconium(IV) isopropoxide/isopropanol**

At room temperature, 10.0 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole and 1.05 g of (+)-L-tartaric acid bis-(N-pyrrolidinamide) (0.15 eq.) are suspended in 72 ml of methyl isobutyl ketone. The suspension is heated at 40-45°C and 12 ml of MIBK are distilled off for azeotropic removal of water present in the mixture. At this temperature, 0.53 g of zirconium(IV) isopropoxide/isopropanol (0.05 eq.) is added and stirring is continued for 1 hour. After cooling to 30°C, 0.16 ml of N-ethyl-diisopropylamine is added. 5.5 g of cumene hydroperoxide (80% in cumene) are then slowly metered in. Stirring is continued at 30°C until the exothermic oxidation process has ended (20 hours, monitored by TLC or HPLC). HPLC of the reaction shows 82% of title compound in an optical purity of > 95%.

**9. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole  
[= (-)-pantoprazole or (S)-pantoprazole] with catalytic amounts of (+)-L-tartaric acid bis-  
(N-pyrrolidinamide) and zirconium(IV) n-propoxide**

At room temperature, 50.0 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole and 13.9 g of (+)-L-tartaric acid bis-(N-pyrrolidinamide) (0.40 eq.) are suspended in 360 ml of methyl isobutyl ketone. The suspension is heated at 40-45°C and 60 ml of MIBK are distilled off for azeotropic removal of water present in the mixture. At this temperature, 6.4 g of zirconium(IV) n-propoxide (70% in propanol, 0.10 eq.) are added, and stirring is continued for 1 hour. After cooling to 30°C, 1.8 ml of N-ethyl-diisopropylamine are added. 27.1 g of cumene hydroperoxide (80% in cumene) are then slowly metered in. Stirring is continued at 30°C until the exothermic oxidation process has ended (20 hours, monitored by TLC or HPLC: chemical purity: 90% of pantoprazole sulfoxide). The suspension is diluted with 120 ml of 2-propanol and quenched with 1.69 g of sodium thiosulphate in 100 ml of saturated sodium bicarbonate solution and stirred for at least 2 hours. After phase separation, the mixture is washed twice with 50 ml of saturated sodium bicarbonate solution. 350 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to 12.5-13 using 10 ml of aqueous sodium hydroxide solution (40% (w/w)). After phase separation, the methyl isobutyl ketone

phase is extracted 2 more times with 100 ml of water and 2 ml of aqueous sodium hydroxide solution (40% (w/w)) at pH = 12.5-13. The combined aqueous phases are reextracted twice with 50 ml of methyl isobutyl ketone and subjected to incipient distillation at 40°C under reduced pressure. At 40-45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH = 9.0. Under pH control, stirring is continued for another 12 hours. The beige crystals are filtered off and washed twice with in each case 50 ml of water.

This gives the title compound in a yield of 85% of theory in a chemical purity of 95% and an optical purity of > 95%. To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and again precipitated at pH = 9.0 using acetic acid (10%). This gives the title compound in a yield of 75-80% of theory in a chemical purity of > 98% and an optical purity of > 99%.

**10. (-)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphonyl]-1H-benzimidazole [ = (-)-pantoprazole or (S)-pantoprazole] with (+)-L-tartaric acid bis-(N,N-pyrrolidinamide) and hafnium(IV) tert-butoxide**

At room temperature, 3.67 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole, 4.10 g of (+)-L-tartaric acid bis-(N,N-pyrrolidinamide) and 2.60 ml of hafnium(IV) tert-butoxide are suspended in 18.5 ml of methyl isobutyl ketone. The mixture is heated at 40°C for 1 hour, during which an almost clear solution is formed. After cooling to room temperature, 0.74 ml of N-ethyl-diisopropylamine is added. 2.2 ml of cumene hydroperoxide are then slowly metered in. Stirring is continued at room temperature until the oxidation process has ended (48 hours, monitored by TLC). The clear solution is diluted with 20 ml of methyl isobutyl ketone and quenched with 0.3 g of sodium thiosulphate in 25 ml of saturated sodium bicarbonate solution and stirred for a further 14 hours. After phase separation, the methyl isobutyl ketone phase is washed two more times with 10 ml of saturated sodium bicarbonate solution. 30 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to 13 using 40% strength (w/w) aqueous sodium hydroxide solution. After phase separation, the methyl isobutyl ketone phase is once more extracted with 10 ml of water at pH = 13. The aqueous phases are combined and, at 40°C and under reduced pressure, subjected to incipient distillation. At 40-45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH 9.0. Under pH control, stirring is continued for another 12 hours. The beige crystals are filtered off and washed with 10 ml of water. This gives the title compound in a yield of 2.5 g (65% of theory) in an optical purity of > 95%. To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and again precipitated at pH = 9.0 using acetic acid (10%).

**11. (+)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole  
[= (+)-pantoprazole or (R)-pantoprazole] with catalytic amounts of (-)-D-tartaric acid bis-  
(N-pyrrolidinamide) and zirconium(IV) n-propoxide**

At room temperature, 50.0 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole and 19.5 g of (-)-D-tartaric acid bis-(N-pyrrolidinamide) (0.56 eq.) are suspended in 360 ml of methyl isobutyl ketone (MIBK). The suspension is heated at 40–45°C and 60 ml of MIBK are distilled off for azeotropic removal of water present in the mixture. At this temperature, 14.0 g of zirconium(IV) n-propoxide (70% in propanol, 0.22 eq.) are added, and the mixture stirring is continued for 1 hour. After cooling to 25°C, 3.5 ml of N-ethyl-diisopropylamine are added. 27.1 g of cumene hydroperoxide (80% in cumene) are then slowly metered in. Stirring is continued at 30°C until the exothermic oxidation process has ended (5–10 hours, monitored by TLC or HPLC). The suspension is diluted with 60 ml of 2-propanol and quenched with 1.69 g of sodium thiosulphate in 100 ml of saturated sodium bicarbonate solution and stirred for at least 2 hours. After phase separation, the mixture is washed twice with 50 ml of saturated sodium bicarbonate solution. 150 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to 12.5–13 using 10 ml of aqueous sodium hydroxide solution (40% (w/w)). After phase separation, the methyl isobutyl ketone phase is extracted again with 75 ml of water and with 5 ml of aqueous sodium hydroxide solution (40% (w/w)) at pH = 13. The mixture is then extracted again with 75 ml of water at pH = 12.5–13. The combined aqueous phases are reextracted with 100 ml of methyl isobutyl ketone and, at 40°C and under reduced pressure, subjected to incipient distillation. At 40–45°C, (-)-pantoprazole is precipitated by addition of 10% strength acetic acid to pH = 9.0. Under pH control, stirring is continued for another 12 hours. The beige crystals are filtered off and washed twice with in each case 50 ml of water.

This gives the title compound in a yield of 80% of theory in a chemical purity of 95% and an optical purity of > 95%. To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and again precipitated at pH = 9.0 using acetic acid (10%). This gives the title compound in a yield of 70% of theory in a chemical purity of > 97% and an optical purity of > 98%.

**12. (+)-5-Difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole  
[= (+)-pantoprazole or (R)-pantoprazole] with catalytic amounts of (-)-D-tartaric acid bis-  
(N-pyrrolidinamide) and zirconium(IV) n-propoxide**

At room temperature, 50.0 g of 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylthio]-1H-benzimidazole and 13.9 g of (-)-D-tartaric acid bis-(N-pyrrolidinamide) (0.40 eq.) are suspended in 360 ml of methyl isobutyl ketone. The suspension is heated at 40–45°C and 60 ml of MIBK are distilled off for azeotropic removal of water present in the mixture. At this temperature, 6.4 g of zirconium(IV) n-propoxide (70% in propanol, 0.10 eq.) are added, and the mixture is stirred for 1 hour. After cooling to 30°C, 1.8 ml of N-ethyl-diisopropylamine are added. 27.1 g of cumene hydroperoxide (80% in cumene) are then slowly metered in. Stirring is continued at 30°C until the exothermic oxidation process has ended (20 hours, monitored by TLC or HPLC). The suspension is diluted with 120 ml of

2-propanol and quenched with 1.69 g of sodium thiosulphate in 100 ml of saturated sodium bicarbonate solution and stirred for at least 2 hours. Further work-up is carried out analogously to Example 11. This gives the title compound in a yield of 85% of theory in an optical purity of > 95%. To increase the purity, (-)-pantoprazole is dissolved in water/aqueous sodium hydroxide solution at pH = 13 and again precipitated at pH = 9.0 using acetic acid (10%). This gives the title compound in a yield of 75% of theory in a chemical purity of > 97% and an optical purity of > 98%.

**13. (S)-5-Methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulphonyl]-1H-benzimidazole Na salt [(S)-omeprazole Na salt] with (+)-L-tartaric acid bis-(N-pyrrolidinamide) and zirconium(IV) n-propoxide**

At room temperature, 1.50 g of 5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylthio]-1H-benzimidazole, 1.87 g of (+)-L-tartaric acid bis-(N-pyrrolidinamide) (1.6 eq.) and 1.37 g of zirconium(IV) n-propoxide (70% in propanol, 0.64 eq.) are suspended in 8.5 ml of methyl isobutyl ketone. The mixture is heated at 40°C for one hour, during which an almost clear solution is formed. After cooling to room temperature, 0.33 ml of N-ethyldiisopropylamine is added. 0.86 ml of cumene hydroperoxide (80% in cumene) are then slowly metered in. Stirring is continued at room temperature until the exothermic oxidation process has ended (5-10 hours, monitored by TLC or HPLC). The suspension is diluted with 5 ml of methyl isobutyl ketone and quenched with 57 mg of sodium thiosulphate and 0.7 g of sodium chloride in 7 ml of saturated sodium bicarbonate solution and stirred for at least 2 hours. After phase separation, the mixture is washed twice with 3 ml of saturated sodium bicarbonate solution. 5 ml of water are added to the methyl isobutyl ketone phase, and the pH is adjusted to 12.5-13 using 0.5 ml of aqueous sodium hydroxide solution (40% (w/w)). After phase separation, the methyl isobutyl ketone phase is extracted 2 more times with in each case 2 ml of water and 0.15 ml of aqueous sodium hydroxide solution (40% (w/w)) at pH = 12.5-13. The combined aqueous phases are distilled at 40°C under reduced pressure. 10 ml of acetonitrile are added to the residue and the mixture is concentrated to half its original volume, giving the product as an oil. Stirring is continued for a further 12 hours, resulting in crystallization of the product. The beige crystals are filtered off. This gives the title compound in a yield of 50% of theory in a chemical purity of 85% and an optical purity of > 95%. Optical rotation  $[\alpha]_D^{20} = +40.0$  (c = 1, water)

**Patent claims**

1. Process for preparing optically pure PPI having a sulphinyl structure in enantiomerically pure or enantiomerically enriched form by oxidation of the corresponding sulphides, characterized in that the oxidation is carried out in the presence of a chiral zirconium complex or a chiral hafnium complex.
2. Process for preparing optically pure PPI having a sulphinyl structure in enantiomerically pure or enantiomerically enriched form by oxidation of the corresponding sulphides, characterized in that the oxidation is carried out in the presence of a chiral zirconium complex.
3. Process according to Claim 1, characterized in that the optically pure PPI having a sulphinyl structure is obtained in an optical purity of > 90%.
4. Process according to Claim 1, characterized in that the oxidation is carried out using cumene hydroperoxide.
5. Process according to Claim 1, characterized in that zirconium(IV) acetylacetonate, zirconium(IV) butoxide, zirconium(IV) tert-butoxide, zirconium(IV) ethoxide, zirconium(IV) n-propoxide, zirconium(IV) isopropoxide or zirconium(IV) isopropoxide/isopropanol complex or hafnium(IV) acetylacetonate, hafnium(IV) butoxide, hafnium(IV) tert-butoxide, hafnium(IV) ethoxide, hafnium(IV) n-propoxide, hafnium(IV) isopropoxide or hafnium(IV) isopropoxide/isopropanol complex is used.
6. Process according to Claim 2, characterized in that zirconium(IV) acetylacetonate, zirconium(IV) butoxide, zirconium(IV) tert-butoxide, zirconium(IV) ethoxide, zirconium(IV) n-propoxide, zirconium(IV) isopropoxide or zirconium(IV) isopropoxide/isopropanol complex is used.
7. Process according to Claim 1, characterized in that the chiral auxiliary used is a chiral tartaric acid derivative.
8. Process according to Claim 1, characterized in that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-diallylamide), (+)-L-tartaric acid bis-(N,N-dibenzylamide), (+)-L-tartaric acid bis-(N,N-diisopropylamide), (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-piperidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), (+)-L-tartaric acid bis-(N-cycloheptylamide), (+)-L-tartaric acid bis-(N-4-methyl-N-piperazinamide), dibutyl (+)-L-tartrate, di-tert-butyl (+)-L-tartrate, diisopropyl (+)-L-tartrate, dimethyl (+)-L-tartrate, diethyl (+)-L-tartrate, (-)-D-tartaric acid bis-(N,N-diallylamide), (-)-D-tartaric acid bis-(N,N-dibenzylamide), (-)-D-tartaric acid bis-(N,N-diisopropylamide), (-)-D-tartaric acid bis-(N,N-dimethylamide), (-)-D-tartaric acid bis-(N-pyrrolidinamide), (-)-D-tartaric acid bis-(N-piperidinamide), (-)-D-tartaric acid bis-(N-morpholinamide), (-)-D-tartaric acid bis-(N-cycloheptylamide), (-)-D-tartaric acid bis-(N-4-methyl-N-

piperazinamide), dibutyl (-)-D-tartrate, di-tert-butyl (-)-D-tartrate, diisopropyl (-)-D-tartrate, dimethyl (-)-D-tartrate or diethyl (-)-D-tartrate.

**9.** Process according to Claim 1, characterized in that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), (-)-D-tartaric acid bis-(N,N-dimethylamide), (-)-D-tartaric acid bis-(N-pyrrolidinamide) or (-)-D-tartaric acid bis-(N-morpholinamide).

**10.** Process according to Claim 1, characterized in that the oxidation is carried out in the presence of an organic base.

**11.** Process according to Claim 1, characterized in that the oxidation is carried out in the presence of a tertiary amine.

**12.** Process according to Claim 1, characterized in that the oxidation is carried out in organic solvents.

**13.** Process according to Claim 1, characterized in that the oxidation is carried out in organic solvents comprising 0 to 0.3% by volume of water.

**14.** Process according to Claim 1, characterized in that the oxidation is carried out in an organic solvent which essentially comprises methyl isobutyl ketone.

**15.** Process according to Claim 1, characterized in that the zirconium component used is zirconium(IV) acetylacetonate, zirconium(IV) butoxide, zirconium(IV) tert-butoxide, zirconium(IV) ethoxide, zirconium(IV) n-propoxide, zirconium(IV) isopropoxide, or zirconium(IV) isopropoxide/isopropanol complex, and that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-diallylamide), (+)-L-tartaric acid bis-(N,N-dibenzylamide), (+)-L-tartaric acid bis-(N,N-diisopropylamide), (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-piperidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), (+)-L-tartaric acid bis-(N-cycloheptylamide), (+)-L-tartaric acid bis-(N-4-methyl-N-piperazinamide), dibutyl (+)-L-tartrate, di-tert-butyl (+)-L-tartrate, diisopropyl (+)-L-tartrate, dimethyl (+)-L-tartrate, diethyl (+)-L-tartrate, (-)-D-tartaric acid bis-(N,N-diallylamide), (-)-D-tartaric acid bis-(N,N-dibenzylamide), (-)-D-tartaric acid bis-(N,N-diisopropylamide), (-)-D-tartaric acid bis-(N,N-dimethylamide), (-)-D-tartaric acid bis-(N-pyrrolidinamide), (-)-D-tartaric acid bis-(N-piperidinamide), (-)-D-tartaric acid bis-(N-morpholinamide), (-)-D-tartaric acid bis-(N-cycloheptylamide), (-)-D-tartaric acid bis-(N-4-methyl-N-piperazinamide), dibutyl (-)-D-tartrate, di-tert-butyl (-)-D-tartrate, diisopropyl (-)-D-tartrate, dimethyl (-)-D-tartrate or diethyl (-)-D-tartrate.

**16.** Process according to Claim 1, characterized in that the zirconium component used is zirconium(IV) acetylacetonate, zirconium(IV) butoxide, zirconium(IV) tert-butoxide, zirconium(IV) ethoxide, zirconium(IV) n-propoxide, zirconium(IV) isopropoxide, or zirconium(IV) isopropoxide/isopropanol complex, that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-diallylamide), (+)-L-tartaric acid bis-

(N,N-dibenzylamide), (+)-L-tartaric acid bis-(N,N-diisopropylamide), (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-piperidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), (+)-L-tartaric acid bis-(N-cycloheptylamide), (+)-L-tartaric acid bis-(N-4-methyl-N-piperazinamide), dibutyl (+)-L-tartrate, di-tert-butyl (+)-L-tartrate, diisopropyl (+)-L-tartrate, dimethyl (+)-L-tartrate, diethyl (+)-L-tartrate, (-)-D-tartaric acid bis-(N,N-diallylamide), (-)-D-tartaric acid bis-(N,N-dibenzylamide), (-)-D-tartaric acid bis-(N,N-diisopropylamide), (-)-D-tartaric acid bis-(N,N-dimethylamide), (-)-D-tartaric acid bis-(N-pyrrolidinamide), (-)-D-tartaric acid bis-(N-piperidinamide), (-)-D-tartaric acid bis-(N-morpholinamide), (-)-D-tartaric acid bis-(N-cycloheptylamide), (-)-D-tartaric acid bis-(N-4-methyl-N-piperazinamide), dibutyl (-)-D-tartrate, di-tert-butyl (-)-D-tartrate, diisopropyl (-)-D-tartrate, dimethyl (-)-D-tartrate or diethyl (-)-D-tartrate and that the oxidation is carried out in the presence of an organic base.

**17.** Process according to Claim 1, characterized in that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide), (+)-L-tartaric acid bis-(N-morpholinamide), (-)-D-tartaric acid bis-(N,N-dimethylamide), (-)-D-tartaric acid bis-(N-pyrrolidinamide) or (-)-D-tartaric acid bis-(N-morpholinamide) and that the oxidation is carried out in the presence of an organic base.

**18.** Process according to Claim 1, characterized in that (S)-5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (S)-5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (S)-2-[3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (S)-2-[4-[3-methoxypropoxy]-3-methylpyridin-2-yl)methylsulphinyl]-1H-benzimidazole, (S)-5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridylmethyl)sulphinyl]-1H-imidazo(4,5-b)pyridine, (R)-5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (R)-5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (R)-2-[3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (R)-2-[(4-(3-methoxypropoxy)-3-methylpyridin-2-yl)methylsulphinyl]-1H-benzimidazole or (R)-5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridylmethyl)sulphinyl]-1H-imidazo(4,5-b)pyridine is prepared by the process.

**19.** Process according to Claim 1, characterized in that the chiral auxiliary used is (-)-D-tartaric acid bis-(N,N-dimethylamide), (-)-D-tartaric acid bis-(N-pyrrolidinamide) or (-)-D-tartaric acid bis-(N-morpholinamide) and that the process product prepared is (+)-pantoprazole.

**20.** Process according to Claim 1, characterized in that the zirconium component used is zirconium(IV) n-propoxide, zirconium(IV) isopropoxide or zirconium(IV) isopropoxide/isopropanol complex, that the chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide) or (+)-L-tartaric acid bis-(N-morpholinamide), that the oxidation is carried out using cumene hydroperoxide and that the process product prepared is (-)-pantoprazole.

**21.** Process according to Claim 1, characterized in that the zirconium component used is zirconium(IV) n-propoxide, zirconium(IV) isopropoxide or zirconium(IV) isopropoxide/isopropanol complex, that the



chiral auxiliary used is (+)-L-tartaric acid bis-(N,N-dimethylamide), (+)-L-tartaric acid bis-(N-pyrrolidinamide) or (+)-L-tartaric acid bis-(N-morpholinamide), that the oxidation is carried out using cumene hydroperoxide in the presence of a tertiary amine and that the process product prepared is (-)-pantoprazole.

**22.** (S)-5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (S)-5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (S)-2-[3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (S)-2-[[4-(3-methoxypropoxy)-3-methylpyridin-2-yl]methylsulphinyl]-1H-benzimidazole or (S)-5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridylmethyl)sulphinyl]-1H-imidazo[4,5-b]pyridine, (R)-5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (R)-5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (R)-2-[3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridinyl)methylsulphinyl]-1H-benzimidazole, (R)-2-[[4-(3-methoxypropoxy)-3-methylpyridin-2-yl]methylsulphinyl]-1H-benzimidazole or (R)-5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridylmethyl)sulphinyl]-1H-imidazo[4,5-b]pyridine prepared by the process according to Claim 1.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/ 03/13605

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07D401/12 C07D471/04

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 02535 A (COTTON HANNA KRISTINA ;LARSSON ERIK MAGNUS (SE); ASTRA AB (SE); SO) 1 February 1996 (1996-02-01) cited in the application claim 1	1-21
X	examples	22
Y	BONCHIO ET AL: "The first Chiral Zirconium(IV) catalyst for highly stereoselective sulfoxidation" JOURNAL OF ORGANIC CHEMISTRY., vol. 64, no. 4, 1999, pages 1326-1330, XP002242676 AMERICAN CHEMICAL SOCIETY. EASTON., US ISSN: 0022-3263 the whole document	1-21

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

17 March 2004

Date of mailing of the international search report

24/03/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

De Jong, B

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/ 03/13605

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	COTTON H. ET AL: "Asymmetric synthesis of esomeprazole" TETRAHEDRON: ASYMMETRY, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 11, no. 18, 22 September 2000 (2000-09-22), pages 3819-3825, XP004224163 ISSN: 0957-4166	1-21
X	examples	22

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/03/13605

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9602535	A	01-02-1996	SE 504459 C2 17-02-1997
			AT 242233 T 15-06-2003
			AU 688074 B2 05-03-1998
			AU 2994895 A 16-02-1996
			BR 9508292 A 23-12-1997
			CA 2193994 A1 01-02-1996
			CN 1157614 A ,B 20-08-1997
			CZ 9700064 A3 11-06-1997
			DE 69530987 D1 10-07-2003
			DK 773940 T3 15-09-2003
			EE 3354 B1 15-02-2001
			EP 0773940 A1 21-05-1997
			FI 970102 A 10-01-1997
			HK 1008331 A1 21-11-2003
			HR 950401 A1 31-10-1997
			HU 76642 A2 28-10-1997
			IL 114477 A 24-07-2001
			JP 10504290 T 28-04-1998
			NO 970153 A 14-01-1997
			NZ 289959 A 26-01-1998
			PL 318165 A1 26-05-1997
			PT 773940 T 31-10-2003
			RU 2157806 C2 20-10-2000
			SE 9402510 A 16-01-1996
			WO 9602535 A1 01-02-1996
			SI 773940 T1 29-02-2004
			SK 4897 A3 06-08-1997
			TR 960063 A2 21-06-1996
			US 5948789 A 07-09-1999
			ZA 9505724 A 15-01-1996